



**Assessment of the Occurrence of Cancer
Chasewood Subdivision, Fort Bend County, TX
2003-2012
March 21, 2016**

Prepared by the
Texas Department of State Health Services

Table of Contents

Executive Summary	1
Background	2
Methods.....	2
Data Sources	3
Statistical Analysis.....	3
Results	4
Table 1.	4
Discussion.....	5
Additional Information	5

Executive Summary

Citizen concern, conveyed by the City of Houston Health Department, prompted the Environmental and Injury Epidemiology and Toxicology Unit (EIET) and Texas Cancer Registry (TCR) of the Texas Department of State Health Services (DSHS) to examine the occurrence of cancer in the Chasewood subdivision located in Fort Bend County, Texas.

DSHS followed agency protocol and the Centers for Disease Control and Prevention (CDC) and Council of State and Territorial Epidemiologists (CSTE) 2013 guidelines to investigate the occurrence of six types of all-ages cancers in a geographic area selected to correspond with community members' concerns. In accordance with these guidelines, the purpose of this assessment was to determine whether the observed number of cancer cases is statistically significantly greater than expected. It was not intended to determine the cause of the observed cancers or identify possible associations with any risk factors.

DSHS staff analyzed TCR data available for a 10-year period spanning from 2003-2012. United States Census data was used to estimate the population in the selected geographic area, which consisted of four census tracts. To evaluate the occurrence of cancer in the area investigated, the number of observed cancer cases was compared to what would be expected for the area based on cancer rates in Texas. Standardized incidence ratios (SIRs) were calculated as the number of observed cases divided by the number of expected cases in the area of concern for the 10-year period (2003-2012). A 95 percent confidence interval (CI) was calculated for each SIR to determine statistical significance.

Observed numbers of three of the cancers were statistically significantly less than expected. The remainder were within the range of what was expected, based on cancer rates in Texas. None of the cancers were statistically significantly higher than expected.

Background

The City of Houston Health Department contacted the Texas Department of State Health Services (DSHS) to convey citizen's concerns regarding cancer in the Chasewood subdivision, located in Ft. Bend County, Texas. Local residents were concerned about cancers in their neighborhood, which they thought could be due to potential exposure to drinking water containing radionuclides at concentrations higher than the Maximum Contaminant Level (MCL) established by the U.S. Environmental Protection Agency (EPA). This prompted the DSHS Environmental and Injury Epidemiology and Toxicology Unit (EIET) and Texas Cancer Registry (TCR) to examine the occurrence of cancer in this community.

The Centers for Disease Control and Prevention (CDC) and Council of State and Territorial Epidemiologists (CSTE) define a cancer cluster as a greater than expected number of cancer cases that occurs within a group of people in a geographic area over a defined period of time¹. DSHS followed the CDC and CSTE 2013 Guidelines for Investigating Suspected Cancer Clusters and Responding to Community Concerns¹ to investigate the occurrence of cancer in this community.

The CDC and CSTE guidelines include four steps¹. The first step is to collect information about the community's concerns. The second step, reported here, is to determine whether the observed number of cancer cases is statistically significantly greater than expected. It is important to note that the data and statistical analysis conducted at this step cannot determine if cancers observed in the community are associated with environmental, lifestyle, or other risk factors.

The guidelines also provide additional steps that can be followed when appropriate. The third step is to evaluate the feasibility of performing an epidemiologic study to examine if exposure to a specific risk factor is associated with the suspected cancer cluster, and the fourth step is to conduct an epidemiologic study, if deemed feasible in step three. Many factors are considered in making the determination to progress to steps three or four. The CDC and CSTE guidelines state, "only a small fraction of cancer cluster inquiries might meet the statistical and etiological criteria to support a cluster investigation through all the steps outlined...."¹

Methods

Consistent with DSHS protocol² and the CDC and CSTE guidelines, DSHS collaborated with the City of Houston Health Department (HHD) to select the geographic area, time frame, and cancers to be included in this analysis in accordance with concerns expressed by the community.

The following all-age cancer types were included in the analysis: liver, pancreas, lung, lymphocytic leukemia (acute and chronic combined), myeloid leukemia (acute and chronic combined), and oropharynx. The HHD also requested that DSHS include esophageal, gallbladder, nasal, and bone cancers. However, because there were five or fewer cases of each of

¹ Centers for Disease Control and Prevention. *Investigating Suspected Cancer Clusters and Responding to Community Concerns*. MMWR, 2013. 62: p. 22.

these cancers in the area of concern during this time frame, they could not be included in the analysis per agency protocol².

DSHS evaluated ten years of available cancer data because the exact years of concern were not known, and the selected timeframe would include more recent cancers that the community was concerned about.

The geographic area investigated was selected in order to encompass all homes in the neighborhood of concern. The four census tracts comprising the area investigated can be seen in Figure 1.

This document outlines the results from step two of the CDC and CSTE guidelines, and only addresses the question, “Is there a statistically significant excess of cancer in the area of investigation?”

Data Sources

For each cancer type, the number of cases observed from *years* in the area included in the investigation was obtained from the TCR (Incidence – Texas, 1995-2012, SEER*Prep 2.5.2). The TCR is responsible for the collection, maintenance, and dissemination of high-quality Texas population-based cancer data, and meets national CDC timeliness and data quality standards, as well as North American Association of Central Cancer Registry certification standards. Cancers were defined according to Site Recode ICD-O-3/WHO 2008 Definitions³. Statewide cancer rates for the same time period were also obtained from the TCR.

Population estimates for 2003-2012 were calculated using linear interpolation based on population counts obtained from the United States Decennial Census⁴ for the years 2000 and 2010. This method, outlined by the United States Census Bureau⁵, assumed population growth occurred in a linear manner.

Statistical Analysis

To determine if a statistically significant excess of cancer existed in the area investigated, the number of observed cancer cases was compared to what would be expected for the area based on cancer rates in Texas. Characteristics such as race, sex, and age are closely related to cancer. To ensure that differences between the numbers of observed and expected cancer cases are not simply due to differences in these demographic characteristics, the expected numbers of cancer

² Texas Department of State Health Services. *Protocol for Responding to Community Cancer Cluster Concerns*. 2016. Available online: <http://www.dshs.state.tx.us/epitox/CancerClusters/Protocol-for-Responding-to-Community-Cancer-Cluster-Concerns.pdf>

³ National Cancer Institute, Surveillance, Epidemiology and End Results Program. *Site Recode ICD-O-3/WHO 2008 Definition*. Available online: http://seer.cancer.gov/siterecode/icdo3_dwhohome/index.html

⁴ United States Census Bureau. *American FactFinder*. 2012; Available from: <http://factfinder2.census.gov/faces/nav/jsf/pages/index.xhtml>

⁵ US Census Bureau. *Methodology for the Intercensal Population and Housing Unit Estimates: 2000 to 2010*. 2012; Available online: http://www.census.gov/popest/methodology/2000-2010_Intercensal_Estimates_Methodology.pdf

cases were calculated by multiplying the age-, sex-, and race-specific cancer incidence rates of Texas residents (reference population) by the number of people in the corresponding demographic groups in the area of investigation.

Standardized incidence ratios (SIRs) were calculated to determine if an excess of cancer exists in the area. The SIR is the number of observed cases compared to (divided by) the number of expected cases for each cancer type. A SIR greater than 1.00 indicates that the observed number of cases of a specific cancer type is higher than expected and a SIR less than 1.00 indicates that the observed number of cases of a specific cancer type is lower than expected.

Few, if any, communities will have exactly the same rate as the average state rate for a similar population; most will be higher or lower. Therefore, 95 percent confidence intervals (CI) were calculated for the SIRs to determine if the observed number of cases was statistically significantly different than expected. If a 95 percent CI (range) includes 1.00, no statistically significant excess (or reduction) of cancer is indicated. If a 95 percent CI does not contain 1.00, the SIR is outside the expected range and is statistically significant. When using a 95 percent CI, 5 percent of SIR values calculated is expected to be statistically significantly higher or lower than the state average due to random chance alone.

In all cases, when results are described as significant or not significant, DSHS is referring only to statistical significance, with the understanding that all cases of cancer are significant to the individual, the family, and friends of the individuals who are affected.

Results

Table 1 presents the number of observed cases, the number of expected cases, the SIRs, and the corresponding 95 percent CIs for each cancer type evaluated in the area of investigation. The number of liver, lung, and oropharyngeal cancers was statistically significantly lower than expected. The number of pancreatic cancers and lymphocytic and myeloid leukemias was not statistically significantly different than expected. None of the cancers analyzed was statistically significantly greater than expected.

Table 1. Standardized Incidence Ratios (SIRs) and 95 percent Confidence Intervals (CIs) for Selected Cancers in Four Ft. Bend County Census Tracts Analyzed Together, 2003–2012.

Cancer Type	Observed	Expected	SIR	95% CI
All Ages				
Liver	9	21.1	0.43	(0.19, 0.81)*
Pancreas	15	19.8	0.76	(0.42, 1.25)
Lung	68	99.5	0.68	(0.53, 0.87)*
Lymphocytic Leukemia	6	7.8	0.77	(0.28, 1.67)
Myeloid Leukemia	6	7.7	0.78	(0.29, 1.70)
Oropharynx	8	15.9	0.50	(0.22, 0.99)*

* Indicates observed number of cancer cases is statistically significantly **lower** than expected

Discussion

Consistent with DSHS protocol, and the second step of the CDC and CSTE guidelines for investigating suspected cancer clusters, the primary purpose of this step (assessment) is to determine whether the observed number of cases is statistically significantly greater than expected¹. It is not intended to determine the cause of the observed cancers or identify possible associations with any risk factors.

The assessment step in a cancer cluster investigation has several inherent limitations, and results should be interpreted with these limitations in mind. Cancer is not a single disease, but rather many different diseases. Different types of cancers vary in etiologies (causes or origins) and may not share the same predisposing factors. Cancers may be associated with a variety of factors such as genetics, lifestyle, and socioeconomic status. Because cancer is common, cases might appear to occur with alarming frequencies within a community even when the number of cases is within the expected rate for the population.

Additionally, cancer incidence data are based on residence at the time of diagnosis. As people move, it becomes more difficult to determine whether living in the area of investigation is associated with an excess of cancers, because residential history is not tracked. Latency (the time period elapsed between exposure and illness onset) adds to the complexity of this step in the investigation. For most adult cancers, a period of 10 to 40 years can elapse between the beginning of an exposure to a cancer-causing agent and the development of a clinically diagnosable case of cancer. It is possible that former residents who developed cancer no longer lived in the area at the time of diagnosis, and these cases would not be included in this assessment. It is also possible that new people have moved into the area and then were diagnosed with cancer; these cases are included in this assessment.

Conclusion

This assessment did not identify any statistically significant excess of lymphocytic or myeloid leukemia, liver, lung, pancreatic, or oropharynx cancer the area investigated. Community members may be concerned about cancers diagnosed more recently than 2012. However, TCR data are not yet complete for that time period, and could not be included in this analysis. Therefore, DSHS will update this analysis upon request when new data becomes available.

Additional Information

For additional information about cancer clusters, visit the Centers for Disease Control and Prevention, “About Cancer Clusters,” web page at <http://www.cdc.gov/nceh/clusters/about.htm>.

For additional information on cancer risk factors, visit the American Cancer Society, “What Causes Cancer?” web page at <http://www.cancer.org/cancer/cancercauses/index>.

Questions or comments regarding this investigation may be directed to Emily Hall, MPH, Epidemiologist, Environmental & Injury Epidemiology & Toxicology Unit, at 512-776-3723 (email: emily.hall@dshs.state.tx.us) or to Leticia Nogueira, PhD, MPH, Epidemiology Manager, Cancer Epidemiology and Surveillance Branch at 512-776-3422 (email: leticia.nogueira@dshs.state.tx.us).

Figure 1. Selected Census Tracts (2010) for the Chasewood Subdivision and Surrounding Area.

